

UNCLASSIFIED

AD NUMBER
ADB240188
NEW LIMITATION CHANGE
TO Approved for public release, distribution unlimited
FROM Distribution authorized to U.S. Gov't. agencies only; Proprietary Info; Jul 98 Other requests shall be referred to USAMRMC, Fort Detrick, MD 21702-5012
AUTHORITY
USAMRMC ltr, 1 Jun 2001.

THIS PAGE IS UNCLASSIFIED

AD _____

GRANT NUMBER DAMD17-96-1-6102

TITLE: TGF-alpha Expression During Breast Tumorigenesis

PRINCIPAL INVESTIGATOR: Kevin Taylor

CONTRACTING ORGANIZATION: University of Cincinnati
Cincinnati, Ohio 45267-0529

REPORT DATE: July 1998

TYPE OF REPORT: Annual

PREPARED FOR: U.S. Army Medical Research and Materiel Command
Fort Detrick, Maryland 21702-5012

DISTRIBUTION STATEMENT: Distribution authorized to U.S. Government agencies only (proprietary information, Jul 98). Other requests for this document shall be referred to U.S. Army Medical Research and Materiel Command, 504 Scott Street, Fort Detrick, Maryland 21702-5012.

The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or decision unless so designated by other documentation.

**Reproduced From
Best Available Copy**

DTIC QUALITY INSPECTED 4

REPORT DOCUMENTATION PAGE

Form Approved
OMB No. 0704-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.

1. AGENCY USE ONLY (Leave blank)		2. REPORT DATE July 1998	3. REPORT TYPE AND DATES COVERED Annual (1 Jul 97 - 30 Jun 98)	
4. TITLE AND SUBTITLE TGF-alpha Expression During Breast Tumorigenesis			5. FUNDING NUMBERS DAMD17-96-1-6102	
6. AUTHOR(S) Kevin Taylor				
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) University of Cincinnati Cincinnati, Ohio 45267-0529			8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) Commander U.S. Army Medical Research and Materiel Command Fort Detrick, Frederick, Maryland 21702-5012			10. SPONSORING/MONITORING AGENCY REPORT NUMBER	
11. SUPPLEMENTARY NOTES				
12a. DISTRIBUTION / AVAILABILITY STATEMENT Distribution authorized to U.S. Government agencies only (proprietary information, Jul 98). Other requests for this document shall be referred to U.S. Army Medical Research and Materiel Command, 504 Scott Street, Fort Detrick, Maryland 21702-5012.			12b. DISTRIBUTION CODE	
13. ABSTRACT (Maximum 200) Transforming Growth Factor-alpha (TGFα) is an oncogenic growth factor that drives estrogen-independent breast tumor proliferation via an autocrine loop. It's receptor, Epidermal Growth Factor Receptor (EGFR), is a proto-oncogene that may be overexpressed in some breast tumors. Normal breast TGFα and EGFR are expressed in an estrogen-dependent cell specific manner. To determine whether these oncogenes are important in the early stages of tumorigenesis, we compared the TGFα mRNA and protein expression in normal epithelium and tumor cell lines and examined the expression of TGFα and EGFR mRNA and protein expression in normal epithelium, proliferative breast disease, and carcinoma in situ. Breast cancer cell lines expressed primarily the expected 4.6 kb message. No evidence of splice variants was detected. Interestingly, protein expression from freshly isolated normal and tumor epithelium expressed primarily the soluble form of TGFα while the cell lines expressed the transmembrane form. This suggests that tissue culture may affect the normal regulation of this protein. TGFα protein expression in archival human tissues was constant during progression from benign to invasive disease. However, EGFR expression increased at the stage of proliferative breast disease, suggesting that increased EGFR expression is an important early event in the progression towards invasive disease.				
14. SUBJECT TERMS Breast Cancer EGFR TGF-alpha			Breast Tumorigenesis oncogenes autocrine/paracrine	
15. NUMBER OF PAGES 24			16. PRICE CODE	
17. SECURITY CLASSIFICATION OF REPORT Unclassified	18. SECURITY CLASSIFICATION OF THIS PAGE Unclassified	19. SECURITY CLASSIFICATION OF ABSTRACT Unclassified	20. LIMITATION OF ABSTRACT Limited	

19981210 089

FOREWORD

Opinions, interpretations, conclusions and recommendations are those of the author and are not necessarily endorsed by the U.S. Army.

____ Where copyrighted material is quoted, permission has been obtained to use such material.

____ Where material from documents designated for limited distribution is quoted, permission has been obtained to use the material.

____ Citations of commercial organizations and trade names in this report do not constitute an official Department of Army endorsement or approval of the products or services of these organizations.

____ In conducting research using animals, the investigator(s) adhered to the "Guide for the Care and Use of Laboratory Animals," prepared by the Committee on Care and use of Laboratory Animals of the Institute of Laboratory Resources, national Research Council (NIH Publication No. 86-23, Revised 1985).

~~____~~ For the protection of human subjects, the investigator(s) adhered to policies of applicable Federal Law 45 CFR 46.

____ In conducting research utilizing recombinant DNA technology, the investigator(s) adhered to current guidelines promulgated by the National Institutes of Health.

____ In the conduct of research utilizing recombinant DNA, the investigator(s) adhered to the NIH Guidelines for Research Involving Recombinant DNA Molecules.

____ In the conduct of research involving hazardous organisms, the investigator(s) adhered to the CDC-NIH Guide for Biosafety in Microbiological and Biomedical Laboratories.


PI - Signature

7/29/88
Date

Year Two of Predoctoral Award

TABLE OF CONTENTS

Front Cover	1
Standard Form (SF) 298	2
Foreword	3
Table of Contents	4
Introduction	5
Results	9
Conclusions	12
Materials & Methods	14
Bibliographies	15
Figures	21

INTRODUCTION

Breast Cancer As a Process. The etiology of breast cancer can be defined in terms of the genetic changes which predispose breast epithelium to disregulated growth. These genetic changes may be inherited or the result of environmental insults. Regardless of the mechanism, these genetic changes result in the increased expression of oncogenes and the loss of expression of tumor suppressor genes, leading to uncontrolled proliferation (1-3). Endogenous estrogens are among the environmental influences which contribute to breast tumor development. Estrogens play a significant role in the normal growth and development of the breast (4). However, there is now little doubt that estrogenic stimulation of normal breast epithelial proliferation is central to the etiology of breast cancer development (5). Without estrogen, women do not get breast cancer. The mechanism by which estrogen-induced proliferation predisposes to specific genetic changes is not known. Furthermore, the mechanism of estrogen interaction with proto-oncogene products in normal breast tissue has not been extensively examined. This project focuses on the interaction of estrogen and one such proto-oncogene, transforming growth factor α (TGF α), during tumor development.

TGF α in Breast Tumorigenesis. Transforming Growth Factor α is commonly expressed at high levels in invasive breast cancer (6-8). This protein is a dominant oncogene, i.e. when overexpressed in breast epithelium, it produces a fully transformed cell (7-11). However, TGF α is also expressed at high levels in normal breast epithelium, and its synthesis is driven by estrogen (12-17). The function of TGF α in normal epithelium, therefore, appears to be quite different than that in tumor cells. Explaining this dichotomy is one of our goals.

Many oncogenes encode growth factors, therefore the effects of growth factors in malignant cells have been well characterized (18). For instance, we know that the effects of growth factors (including TGF α) are dependent on the differentiated state of the cell, that is a growth factor which causes proliferation at one state of cellular differentiation may induce quiescence in a cell in another state (19). TGF α is expressed in normal epithelium, during pre-malignant proliferation, and in both in situ and invasive carcinoma (8, 20-21). Yet its function is clearly

different over this progression to tumor formation. Understanding the role TGF α plays in normal and premalignant breast tissue is critical to understanding the mechanism through which TGF α mediates its effects in tumors.

Breast Histology. To understand the function of TGF α in the breast, one must understand normal breast histology. Breast epithelium is composed of several epithelial cell types (22). Ducts are lined by luminal and myoepithelial cells both of which are derived from the same stem cell (23-25). The milk-forming unit, or the lobule of the breast, is also composed of two epithelial cell types, alveolar cells and myoepithelial cells (22). The lobular myoepithelial cells are morphologically similar to the ductal myoepithelial cells; however, biochemically they display some differences (our observations). For the most part, myoepithelial differentiation is rare in invasive carcinoma and when ducts are filled with carcinoma at a stage called "in situ carcinoma", the myoepithelial cells are lost, or at best, greatly reduced in number (23). This is true regardless of whether the carcinoma has a more ductal or lobular phenotype. In either case, the cell of origin is thought to reside in the terminal ducto-lobular unit. Hormone-sensitive connective tissue composed of fibroblasts and myofibroblasts surrounds the epithelial cells of the breast. The role these cells play in the process of carcinogenesis is completely unknown. Each of these cell types expresses proto-oncogenes during the normal development and maturation of the breast. The regulation of this expression is the process which goes awry in tumorigenesis. To understand this regulation, we must examine the cell-specific expression of these proto-oncogenes and define the characteristics of normal breast which participate in this regulation.

Our preliminary data suggests that TGF α and its receptor (epidermal growth factor receptor or EGFR) function by two different mechanisms in normal breast. Here we show that the luminal cells of normal, quiescent breast epithelium express TGF α , but not EGFR; therefore, unlike tumor cells, the luminal cells can not respond to secreted TGF α in an autocrine fashion. Myoepithelial cells and adjacent stromal cells express EGFR normally; therefore, in normal breast the target for TGF α production is the myoepithelium and connective tissue. The cellular responses to this

stimulation by TGF α are not known. Between quiescent epithelium and invasive tumor lie intermediate states of proliferation/transformation. Proliferative breast diseases of various morphologic types carry an increased risk of carcinoma and are considered "pre-malignant" phenotypes (27). In the literature and in our own experience, luminal EGFR expression is associated with some proliferative and cystic breast diseases (28). In particular, luminal cells with EGFR also express the estrogen receptor (ER) (28). Thus, alterations in EGFR synthesis may appear very early in proliferating breast, driven by estrogen. In fact, antisense studies with TGF α suggest that induction of TGF α synthesis may be the main mechanism for estrogen-induced proliferation in this cell type (15). By the time breast epithelium has the characteristics of in situ carcinoma, TGF α and EGFR clearly co-localize in the luminal cells. The luminal cells up regulate EGFR and in the fully transformed state, respond to TGF α in an autocrine loop. At this stage there is an inverse correlation between EGFR and ER expression, consistent with the concept that over-stimulation of the TGF α /EGFR pathway leads to estrogen-independent growth (29). Some have suggested that cell transformation by EGFR is the result of a "threshold" level of EGFR expression (10). In this case, one would expect quantitatively increased expression in in situ carcinoma or invasive disease relative to these "pre-malignant" lesions. While the luminal cell is being driven to proliferate, the myoepithelial cell is lost during the transition from pre-malignant proliferative growth to in situ carcinoma. The mechanism by which these cells are lost, and the consequences of their loss are not known.

In this project we explore the role of TGF α in normal and proliferative breast tissue. Understanding the normal role of TGF α in tumor production is critical for the following reasons. First, TGF α is a major estrogen-responsive gene (33); therefore, to understand estrogen-promotion of tumorigenesis, one must know the consequence of estrogen-induced gene expression. Second, TGF α function changes during the process of tumorigenesis. In normal tissue TGF α operates in a paracrine fashion and may be critical for maintaining normal lobular growth. In fact, there is now evidence that lactation is regulated by TGF α (34). However, in tumors TGF α leads to estrogen-independent growth (9). The effects of TGF α in the steps in between, from normal

cell growth to invasive disease, have not been studied. Finally, current experimental therapies target various points in the TGF α signal transduction pathway. For instance, TGF α -EGFR stimulates cells through the MAP kinase pathway (35). This signal transduction pathway involves the proto-oncogene, ras, as well as two kinases which have been well characterized, protein kinase C (PKC) and protein kinase A (PKA) (36). The action of PKC is of particular note, since when EGFR is activated, it results in PKC activation (37-39). As part of an inhibitory feedback loop, PKC then phosphorylates and inhibits the action of EGFR, preventing over stimulation of the cell (40). These are exactly the types of pathways which one would predict are destroyed during the progression to malignancy. Current clinical trials include the use of Bryostatins, an agent which inhibits PKC (41). Given that TGF α can induce proliferation or differentiation, based on the differentiation state of the receptor cell, it is likely that the signaling pathways downstream of EGFR change during tumorigenesis. Therefore, it is critical that we understand the role TGF α is playing in specific breast cells at the time therapies affecting these pathways are instituted.

In this work, we build on our previous work, examining oncogene expression during tumor development and metastasis. Here we shall define the cell-specific expression of TGF α and EGFR in histologically similar premalignant states from women who do or do not progress to invasive tumor. Expression will be correlated with histopathologic types of proliferative breast disease, extent of disease types appearing concurrently, and estrogen receptor status. The observation that TGF α and EGFR must form a paracrine loop between myoepithelial and luminal cells is the result of careful evaluation of benign breast tissues for oncogene expression during a study comparing oncogene expression in benign, invasive, and metastatic tumors (43). In addition, the PI's mentor has collected an extensive set of premalignant breast tissue which are used for these studies. The results of this study will provide an invaluable database of cell-specific gene expression in vivo which can then be correlated with in vitro tissue culture work.

RESULTS

1. Prepare riboprobes for in situ hybridization: subcloning of cDNAs, sequencing of constructs. (Month 0-6).

Conclusion: 925 basepairs of the TGF α gene has been subcloned into the PGEM 3Z plasmid and is in the 5' to 3' orientation relative to the T7 promoter. This goal is completed and presented in 1996-1997 Annual Report.

2. Identify TGF α splice variants in breast cancer cell lines. (Month 6-9).

Initially, the riboprobe was labeled using the Genius Nonradioactive Nucleic Acid Labeling and Detection System (Boehringer-Mannheim Corp., Indianapolis, IN). Due to the poor sensitivity of the digoxigenin-labeled riboprobe labeling was switched from a non-radioactive to a radioactive labeling method. The TGF α probe is now labeled using a standard Pharmacia oligonucleotide labeling protocol (Pharmacia, Piscataway, PA). Seven breast cancer cell lines: Hs-578 Bst, SKBR3, MCF-7, A431, T-47-D, MDA-MB-453, MDA-MB-468, and MCF-10A (ATCC, Rockville, MD) have been examined and show that in all cases the primary mRNA transcript is at 4.6 kb (Figure 1).

Conclusion: Northern blots of isolated RNA on breast cancer cell lines were performed, and probed with the radiolabeled TGF α oligo-probe. All cell lines express the expected 4.6 kb transcript as quoted in the literature (6, 8, 9, 43). No unusual transcripts which would indicate splice variants were detected.

3. Identify TGF α protein sizes in breast cell lines: Western blot. (Month 6-9).

Conclusion: Protein was isolated from: Hs-578 Bst, SKBR3, MCF-7, A431, T-47-D, MDA-MB-453, MDA-MB-468, and MCF-10A cell lines and all primarily express the transmembrane form. This goal is completed and presented in 1996-1997 Annual Report.

4. Repeat 2 and 3 in normal breast cells or organoids. (Month 9-12).

We have examined freshly isolated RNA from eight organoid samples under multiple conditions and have concluded that the TGF α mRNA is too low to detect by Northern blot analysis. Therefore, in order to detect TGF α mRNA expression we are currently developing competitive polymerase chain reaction (PCR).

Previously, Westerns were run on seven breast cancer cell lines, cultured fibroblasts and a fresh breast tissue sample. The fresh breast tissue sample expressed primarily the soluble form (6 kD) of TGF α (reported in 1996-1997 Annual Report). We needed to further investigate if this difference in TGF α protein expression is due to culturing of the cells or an underlying difference between transformed vs. non-transformed epithelial cells. To accomplish this a Western blot was performed on five more patient samples (fresh tissue or isolated organoids) and the cell line MCF-10A. Of the five patient samples one sample was from freshly isolated tumor, and this sample along with the other four normal tissue samples express primarily the soluble form (6 kD) of TGF α (Figure 2). These data indicate that the difference in expression from the fresh tissue samples and cell lines is likely due to culturing of cell lines.

Conclusion: TGF α mRNA from normal epithelium is too low to detect by Northern blot analysis and we are currently in the process of developing competitive PCR to detect message expression. Western blot analysis of patient samples (whole tissue, isolated organoids, and tumor) primarily express the soluble form of TGF α . This is in contrast to cell lines expressing primarily the transmembrane form of the protein, suggesting that fresh tissue may be a more appropriate model to study TGF α protein regulation in breast tissue.

5. Develop splice-specific probes. (Month 9-12).

Conclusion: Northern blot analysis of seven breast cancer cell lines (Figure 1) and fresh breast tissue samples and organoids (Figure 2) using a radiolabeled TGF α oligo-probe do not indicate any potential splice variants.

6. Examine normal and proliferative breast tissue for cell-specific expression of TGF α . (Month 12-24).

Originally most tissues were to be tested for TGF α mRNA expression by in situ hybridization. However, despite good in situ results using cells lines, very inconsistent results were obtained in human archival tissue. To expedite our analysis of TGF α /EGFR expression we identified TGF α expression by immunohistochemistry (IHC). Paraffin embedded formalin fixed tissue from 90 patients were collected and characterized. Protein expression was determined in the luminal epithelial cells and myoepithelial cells of ducts and lobules, as well as the microvasculature, stromal cells and infiltrating lymphocytes. Expression of these proteins in each histological subtype was compared with one another as breast tissue progressed towards invasive disease (i.e. proliferative breast disease, carcinoma in situ (cis), and invasive disease). Our data indicate that TGF α expression does not increase in the luminal epithelial cells of the ducts or lobules as the epithelium progresses through these histopathological stages (Figure 3). In the microvasculature a trend is seen in TGF α upregulation in endothelium adjacent to breast lesions. However, EGFR expression increases in the luminal epithelial cells during the proliferative phase of breast tumorigenesis ($p < 0.006$) compared to normal epithelium, suggesting that upregulation of EGFR is important early on in the progression towards invasive disease (Figure 3). Subsequent to the proliferative stage of progression, EGFR actually decreases.

Conclusion: Analysis of TGF α and EGFR protein expression by IHC in human archival breast tissue reveals: 1. TGF α protein expression is unaltered in the luminal cells as the epithelium progresses through the histopathological subtypes. 2. TGF α is upregulated in endothelium adjacent to breast lesions. 3. EGFR protein expression increases in the luminal cells during the proliferative phase of breast tumorigenesis. This suggests EGFR is an important early event in the progression towards invasive disease.

7. Examine tissue as above for proliferation, estrogen receptor expression, and apoptosis. (Month 18-24).

Estrogen receptor (ER) and progesterone receptor (PR) protein expression were analyzed in a subset of patients. Although there was no consistent difference in ER/PR expression during disease progression, in general high,

functional ER expression correlated with high TGF α expression and low or non-functional ER expression correlated with low TGF α expression (Functional ER expression is defined as the co-expression of PR). These results are consistent with in vitro data indicating TGF α expression can be regulated by an estrogen dependent mechanism (14-16, 33, 43). IHC using the mib-1 monoclonal antibody to determine proliferation has been done and the slide are currently being analyzed. Histological analysis of the human archival breast tissue does not reveal an appreciable amount of apoptotic bodies. Therefore, my thesis committee felt that performing the TUNNELL assay to detect cells undergoing apoptosis would not be informative.

Conclusion: Functional expression of ER (i.e. ER/PR positive) correlated with expression levels of TGF α . IHC using the mib-1 monoclonal antibody to determine proliferation has been performed on all cases and are currently being analyzed. Histological analysis of the human archival breast tissue does not reveal an appreciable amount of apoptotic bodies.

8. Prepare riboprobes for PKC. (Month 24-30).
See below.

9. Examine tissues as above for alterations in PKC isozyme expression. (Month 30-36).
See below.

CONCLUSION

We have subcloned 925 basepairs of the TGF α gene into the PGEM 3Z plasmid and have made a ³²P-labeled probe. Northern blots hybridized with the radiolabeled TGF α probe identified the 4.8 kb mRNA transcript as the primary TGF α message in seven breast cancer cell lines: Hs-578 Bst, SKBR3, MCF-7, T-47-D, MDA-MB-453, MDA-MB-468, and MCF-10A. No splice variants were detected, despite extensive exposure on phosphoimaging plates (Figure 1). Northern blots on 8 patient samples (fresh tissue or organoids) were performed; however, no TGF α mRNA was detected. Using the radiolabeled TGF α probe no mRNA is detected. Further work will focus on developing competitive PCR analysis for TGF α to detect mRNA expression.

Previously, protein was isolated from the same seven breast cancer cell lines (see above) and Westerns performed. The

results indicate that the cell lines express primarily the transmembrane bound form (approximately 22 kD) of TGF α . Western blots performed on patient samples (fresh tissue or organoids) express primarily the soluble form (6 kD) of TGF α . Protein isolated from a tumor specimen also expresses primarily the soluble form of TGF α (Figure 2). These data suggest that the difference in protein expression in the cell lines is due to a tissue culture artefact. Suggesting that fresh tissue may be a more appropriate model to study alterations in TGF α expression during breast tumorigenesis. Immunohistochemical analysis of TGF α /EGFR during progression in individual women indicates that EGFR may play a role in the early proliferative process.

Earlier this year these data were evaluated by my thesis committee who felt that the TGF α data obtained from breast cell lines may not reflect the role of TGF α in the early proliferative process. Therefore, upon their recommendation, I have developed procedures for isolating normal human breast epithelium (organoids) and characterized culture conditions which allow multiple passages of the epithelium in vitro. Furthermore, I have developed methods to characterize the isolated cell types using monoclonal antibodies against keratin 19 (K19) as a luminal marker and keratin 14 (K14) as a myoepithelial marker. Examples of immunohistochemical staining for these proteins in isolated organoids are shown (Figure 4). Having developed these procedures I can now address the following issues: 1. Confirm that TGF α processing differs between in vivo and in vitro by examining TGF α from fresh tissue vs. primary and early passage cultures from the same patient. 2. Determine the role of stromal cells in TGF α protein processing during co-culture of isolated organoids in vitro. 3. Determine whether EGFR expression is altered with conversion from a relatively quiescent organoid to a rapidly proliferating monolayer culture. 4. Investigate the proliferative effect of TGF α and EGF treatment on isolated organoids by immunologic detection of bromo-deoxyuridine (BrDU) incorporation. Given our findings to date, my committee believes that these experiments will allow me to develop functional assays to assess the role of TGF α /EGFR in breast proliferation and develop models in which ER regulation of EGFR expression can be studied. The final objectives of

this proposal were to evaluate PKC isozyme expression. We originally planned to use RNA for isozyme detection. However, over the past two years numerous excellent isozyme-specific antibodies have been developed and are now commercially available. Isozyme expression of matched normal and tumor samples will be analyzed by Western.

MATERIALS & METHODS

Tumor cell culture: Breast tumor cell lines Hs-578 Bst, SKBR3, MCF-7, A431, T-47-D, MDA-MB-453, MDA-MB-468, and MCF-10A were obtained from ATCC and cultured according to their protocols. Cells are passaged weekly using trypsin-EDTA.

Organoid isolation: Breast epithelium (organoids) is dissected from both mastectomy and reduction mammoplasty specimens, minced, and enzymatically digested with collagenase as previously described (45). Briefly, tissue is obtained within 1 hour of surgery, minced with opposing scalpels to 1 mm³ pieces, and enzymatically digested for 12 to 18 hours at 37 °C in a rotary shaker (60 rpm) with 150 U/ml Collagenase Type I (Gibco), 100 U/ml Hyaluronidase (Gibco) in CDM3 media. CDM3 media consists of DMEM/F12 (1:1), 2.6 ng/ml selenium, 100 ng/ml EGF, 0.1 µg/ml fibronectin (all from Collaborative Research, Bedford, MA), 3 µg/ml insulin, 25 µg/ml transferrin, 10⁻¹⁰ M estradiol, 10⁻⁶ M hydrocortisone, 10⁻⁸ M cAMP, 10⁻⁸ M triiodothyronine, 10⁻⁴ M ethanolamine, 10⁻⁴ M phosphoethanolamine, 0.1% BSA, 10 µg/ml ascorbic acid, 20 µg/ml fetuin (all from Sigma, St. Louis, MO), and 1X trace element mix (Biofluids, Rockville, MD). Following digestion, epithelial organoids and fibroblasts are separated from vascular fragments and from one another by sedimentation.

TGFα radiolabeled probe. The probe was labeled with ³²P-CTP using a standard oligonucleotide labeling kit (Pharmacia, Piscataway, PA). Briefly, 25 ng of TGFα plasmid DNA is labeled with ³²P-CTP at 37 °C for 1 hour in a reaction mixture containing random hexanucleotide primers and 50U of Klenow fragment. The labeled probe is then isolated on a Sepharose G50 spin column (Pharmacia, Piscataway, PA).

Northern Blot Analysis. Total cellular RNA is isolated using RNazol (CINNA/BIOTEX) and electrophoresed on formaldehyde-denatured agarose gels as in (65). RNA is transferred onto Nytran for hybridization.

Prehybridization of the filters is carried out for 4-18 hours at 37°C in 50% formamide, 5x SSC, 0.1%

polyvinylpyrrolidone, 0.1% Ficoll, 0.1 mg/ml salmon sperm DNA, 20 mM sodium phosphate (pH 6.8), and 0.1% SDS. For hybridization the appropriate 5'-end labeled oligonucleotide or ³²P-labelled riboprobe is added and incubated at 37° overnight. The filters are then washed in 2x SSC and 1% SDS twice at room temperature, 2x SSC and 1% SDS twice at 37°C, and 0.2x SSC and 1% SDS for 15 minutes at 37°C.

Western blot: Cells or tissues are homogenized in 20mM Tris-HCl, pH 7.4, 2mM EDTA, 0.5mM EGTA for protein analysis (Bradford assay) and then electrophoresed under reducing conditions on 10% acrylamide SDS-PAGE (Laemmli) and transferred to PVDF membranes (Millipore) by the Towbin method. Membranes from each well are cut for replicate analysis, blocked in 5% non-fat dry milk, 0.5M NaCl, 10 mM Tris-HCl, pH 7.5. Primary antibodies (5-10 µg/ml) are incubated in blocking solution at RT overnight. The membranes are washed and incubated one hour at RT with biotinylated anti-mouse/rabbit Ig (1:100, Sigma), washed, and incubated another hour with Streptavidin-alkaline phosphatase (1:200, Sigma) in 5% non-fat milk, 150 mM NaCl, 50 mM Tris-HCl, pH 7.5. The reaction is developed with BCIP/NBT substrate. Samples are calibrated for equal loading by using antibodies against actin.

Immunohistochemistry: 4µm tissue sections are deparaffinized and stained using the Ventanna 320 automated immunostainer (Ventanna Medical Systems, Tucson, AZ). The TGFα antibody Clone 31G7 (Zymed, San Francisco, CA) is used at a dilution of 1:10. The EGFR antibody Clone 213-4.4 (Oncogene Research Products, Cambridge, MA) is used at a dilution of 1:25. The Ki-67 monoclonal antibody mib-1 (Immunotech, West Brook, ME) is used at a dilution of 1:40. The monoclonal antibodies are pretreated with trypsin on the immunostaining system. Each reaction utilizes a horse-radish peroxidase-conjugated secondaries. Slides are then counterstained with hemotoxylin.

BIBLIOGRAPHY

1. Lee EY-HP, Tumor suppressor genes and their alterations in breast cancer. Sem in Cancer Biology. 6:119-125, 1995.
2. Eufjord JE, Thorlacium S, Steinarsdottir M, Valgardsdottir R, Ogmundsdottir HM, Anamthawat-Jonssoon K. p53 abnormalities and genomic instability

- in primary human breast carcinomas. *Cancer Res* 55:646-651, 1995.
3. van Leeuwen F, Nusse R. Oncogene activation and oncogene cooperation in MMTV-induced mouse mammary cancer. *Sem in Cancer Biol.* 6:127-133, 1995.
 4. Richards JE, Shyamala G, Nandi S. Estrogen receptor in normal and neoplastic mouse mammary tissues. *Cancer Res* 34:2764-2772, 1974.
 5. Dickson RB, Lippman ME. Estrogenic regulation of growth and polypeptide growth factor secretion in human breast carcinoma. *Endocrine Rev.* 8:29-43, 1987.
 6. Salomon DS, Kim N, Saeki T, Ciardiello F. Transforming growth factor α : an oncodevelopmental growth factor. *Cancer Cells* 2:389-397, 1990.
 7. Madsen MW, Lyddesfeldt AE, Laursen I, Nielsen KV, Briand P. Altered gene expression of c-myc, epidermal growth factor receptor, transforming growth factor- α , and c-erb-B2 in an immortalized human breast epithelial cell line, HMT-3522, is associated with decreased growth factor requirements. *Cancer Res.* 52:1210-1217, 1992.
 8. Perroteau I, Salomon D, DeBortoli M, Kidwell W, Hazarika P, Pardue R, Dedman J, Tam J. Immunological detection and quantitation of alpha transforming growth factors in human breast carcinoma cells. *Breast Cancer Res and Treat.* 7:201-210, 1986.
 9. Herman ME, Katzenellenbogen BS. Alterations in transforming growth factor- α and - β production and cell responsiveness during the progression of MCF-7 human breast cancer cells to estrogen-autonomous growth. *Cancer Res* 54:5867-5874, 1994.
 10. DiMarco E, Pierce JH, Fleming TP, Kraus MH, Mollay CJ, Aaronson SA, Di Fiore PP. Autocrine interaction between TFG α and the EGF-receptor: quantitative requirements for induction of the malignant phenotype. *Oncogene* 4:831-838, 1989.
 11. Ju WD, Velu TJ, Vass WC, Papageorge AG, Lowry DR. Tumorigenic transformation of NIH 3T3 cells by the autocrine synthesis of transforming growth factor α . *New Biol.* 3:380-388, 1991.
 12. Spitzer E, Zschesche W, Binas B, Grosse R, Erdmann B. EGF and TGF α modulate structural and functional differentiation of the mammary gland from pregnant mice in vitro: possible role of the arachidonic acid pathway. *J Cell Biochem.* 57:495-508, 1995.
 13. Zwiebel JA, Bano M, Nexo E, Salomon DS, Kidwell WR. Partial purification of transforming growth factors from human milk. *Cancer Res.* 46:933-939, 1986.

14. Liu SC, Sanfilippo B, Perroteau I, Derynck R, Salomon DS, Kidwell WR. Expression of transforming growth factor a (TGFa) in differentiated rat mammary tumors: estrogen induction of TGFa production. *Molec Endocrin.* 1:683-692, 1987.
15. Kenney NJ, Saeki T, Gottardis M, Kim N, Garcia-Morales P, Martin MB, Normanno N, Ciardiello F, Day A, Cutler ML, Salomon DS. Expression of transforming growth factor a antisense mRNA inhibits the estrogen-induced production of TGFa and estrogen-induced proliferation of estrogen-responsive human breast cancer cells. *J Cell Physiol.* 156:497-514, 1993.
16. Manni A, Wright C, Gadger B, Bartholomew M, Herlyn M, Mendelsohn, Masui H, Demers L. Role of transforming growth factor-a-related peptides in the autocrine/paracrine control of experimental breast cancer growth in vitro by estradiol, prolactin, and progesterone. *Breast Cancer Res and Treat.* 15:73-83, 1990.
17. Saeki T, Cristiano A, Lynch MJ, Brattain M, Kim N, Normanno N, Denney N, Ciardiello F, Salomon DS. Regulation by estrogen through the 5'-flanking region of the Transforming Growth Factor a gene. *Molec Endocrin.* 5:1955-1963, 1991.
18. Druker BJ, Mamon HJ, Roberts TM. Oncogenes, growth factors, and signal transduction. *New Engl J Med.* 321:1383-1391, 1989.
19. Sauma S, Huang F, Winawer S, Friedman E. Colon goblet cells lose proliferative response to TGFa as they differentiate. *Int J Cancer* 61:848-853, 1995.
20. Snedeker SM, Brown CF, DiAugustine RP. Expression and functional properties of transforming growth factor a and epidermal growth factor during mouse mammary gland ductal morphogenesis. *Proc Natl Acad Sci USA.* 88:276-280, 1991.
21. Scala S, Saeki T, Lynch A, Salomon D, Merino MJ, Bates SE. Coexpression of TGFa, epidermal growth factor receptor, and P-glycoprotein in normal and benign diseased breast tissues. *Diagnostic Molec Pathol.* 4:136-142, 1995.
22. Rudland PS. Histochemical organization and cellular composition of ductal buds in developing human breast: evidence of cytochemical intermediates between epithelial and myoepithelial cells. *J Histochem and Cytochem.* 39:1471-1484, 1991.
23. Rudland PS, Leinster SJ, Winstanley J, Green B, Atkinson M, Zakhour HD. Immunocytochemical

- identification of cell types in benign and malignant breast diseases: variations in cell markers accompany the malignant state. *J Histochem Cytochem.* 41:543-553, 1993.
24. Rudland PS, Barraclough R. Stem cells in mammary gland differentiation and cancer. *J Cell Sci Suppl* 10:95-114, 1988.
 25. Noguchi S, Motomura K, Inaji H, Imaoka S, Koyama H. Clonal analysis of solitary intraductal papilloma of the breast by means of polymerase chain reaction. *Am J Pathol* 144:1320-5, 1994.
 26. Silverstein MJ, Lewinsky BS, Waisman JR, Gierson ED, Colburn WJ, Senofsky GM, Gamagami P. Infiltrating lobular carcinoma: is it different from infiltrating duct carcinoma? *Cancer* 73:1673-1677, 1993
 27. Bodian CA, Perzin KH, Lattes R, Hoffmann P, Abernathy TG. Prognostic significance of benign proliferative breast disease. *Cancer* 71:3896-3907, 1993.
 28. van Agthoven T, Timmermans M, Foekens JA, Dorssers LCJ, Henzen-Logmans SC. Differential expression of estrogen, progesterone, and epidermal growth factor receptors in normal, benign, and malignant human breast tissues using dual staining immunohistochemistry. *Am J Pathol.* 144:1238-1246, 1994.
 29. Newby JC, A'Hern RP, Leek RD, Smith IE, Harris AL, Dowsett M. Immunohistochemical assay for epidermal growth factor receptor on paraffin-embedded sections: validation against ligand-binding assay and clinical relevance in breast cancer. *Br J Cancer* 71:1237-1242, 1995.
 30. Halter SA, Dempsey P, Matsui Y, Stokes MK, Graves-Deal R, Hogan BLM, Coffey RJ. Distinctive patterns of hyperplasia in transgenic mice with mouse mammary tumor virus transforming growth factor- α . *Am J Pathol.* 140:1131-1146, 1992.
 31. Smas CM, Sui HS. Pref-1, a protein containing EGF-like repeats, inhibits adipocyte differentiation. *Cell* 73:725-734, 1993.
 32. Bani D, Riva A, Bigazzi M, Sacchi TB. Differentiation of breast cancer cells in vitro is promoted by the concurrent influence of myoepithelial cells and relaxin. *Br. J. Cancer.* 70:900-904, 1994.
 33. Bates SE, Danidson NE, Valverius EM, Freter CE, Dickson rB, Tam JP, Kudlow JE, Lippman ME, Salomon DS. Expression of transforming growth factor α and its messenger ribonucleic acid in human breast cancer: its

- regulation by estrogen and its possible functional significance. *Molec Endocrinol.* 2:543-555, 1988.
34. Lin CQ, Dempsey PJ, Coffey RJ, Bissell MJ. Extracellular matrix regulates whey acidic protein gene expression by suppression of TGF α in mouse mammary epithelial cells: studies in culture and in transgenic mice. *J Cell Biol.* 129:1115-1126, 1995.
 35. Ahn NG, Seger R, Bratlien RL, Keltz CD, Tonks NK, Krebs EG. Multiple components in an epidermal growth factor-stimulated protein kinase cascade. *J Biol Chem.* 266:4220-4227, 1991.
 36. Buday L, Downward J. Epidermal growth factor regulates p21ras through the formation of a complex of receptor, Grb2, adaptor protein, and Sos nucleotide exchange factor. *Cell* 73:611-620, 1993.
 37. Osborne CK, Hamilton B, Nover M, Ziegler J. Antagonism between epidermal growth factor and phorbol ester tumor promoters in human breast cancer cells. *J Clin Invest.* 67:943-951, 1981.
 38. Fabbro D, Kung W, Roos W, Regazzi R, Eppenberger U. Epidermal growth factor binding and protein kinase C activities in human breast cancer cell lines: possible quantitative relationship. *Cancer Res.* 46:2720-2725, 1986.
 39. Wyss R, Fabbro D, Regazzi R, Borner C, Takahashi A, Eppenberger U. Phorbol ester and epidermal growth factor receptors in human breast cancer. *Anticancer Res.* 7:721-728, 1987.
 40. Schlessinger J. Allosteric regulation of the epidermal growth factor receptor kinase. *J Cell Biol.* 103:2067-2072, 1986.
 41. Blobe GC, Obeid LM, Hannun YA. Regulation of protein kinase C and role in cancer biology. *Cancer and Met Rev.* 13:411-431, 1994.
 42. Lower E, Heffelfinger SC, Miller MA, Fenoglio-Preiser CM. Phosphotyrosine immunoreactivity is a significant prognostic indicator in breast carcinoma patients. *Breast Cancer Res Treat.* 35:277-282, 1995.
 43. Salomon DS, Brandt R, Ciardiello F, Normanno N. Epidermal Growth Factor-related peptides and their receptors in human malignancies. *Critical Review in Oncology/Hematology*, 19:183-232, 1995.
 44. Lan HY, Wei Mu, Yee YY, Nikolic-Patterson DJ, Atkins RC. A simple, reliable, and sensitive method for nonradioactive In Situ Hybridization: Use of Microwave Heating to Improve Hybridization Efficiency and

- Preserve Tissue Morphology. The Journal of
Histochemistry and Cytochemistry 44:281-287, 1996.
45. Petersen OW, vanDeurs B. Preservation of Defined
Phenotypic Traits in Short-Term Cultured Human Breast
Carinoma Derived Epithelial Cells. Cancer Research.
47:856-862, 1987.

Figure 1.

Northern blot showing expression of TGF α mRNA expression using a ^{32}P labeled probe. Lanes are mRNA isolated from cell line: (a) HS-578-Bst, (b) MDA-MB-453, (c) MDA-MB-468, (d) T-47-D, (e) MCF-7, (f) SKBR3, (g) MCF-10A.




Figure 2.

Anti-TGF α Western blot from fresh tissue samples. Lanes are proteins isolated from: (a) MCF-10A cell line, (b) fresh invasive ductal carcinoma, (c) isolated organoids patient A, (d) isolated organoids patient B, (e) fresh breast tissue patient C, (f) fresh breast tissue patient D, (g) commercial 6 kD TGF α .




Figure 3.

Tables showing (a) EGFR protein expression during tumor progression (* $p=0.006$). (b) TGF α protein expression in luminal epithelium during tumor progression. Shown are the mean and standard deviation of immunohistochemical stain intensity for each histologic subtype.







Figure 4.

Immunohistochemistry identifying epithelial cell types in freshly isolated organoids: (a) shows immunoperoxidase staining for cytokeratin K19, identifying luminal epithelial cells (arrow), (b) shows immunoperoxidase staining for cytokeratin 14, identifying myoepithelial cells (arrow). All magnifications are 40x.





DEPARTMENT OF THE ARMY

US ARMY MEDICAL RESEARCH AND MATERIEL COMMAND
504 SCOTT STREET
FORT DETRICK, MARYLAND 21702-5012

REPLY TO
ATTENTION OF:

MCMR-RMI-S (70-1y)

21 JUN 2001

MEMORANDUM FOR Administrator, Defense Technical Information
Center (DTIC-OCA), 8725 John J. Kingman Road, Fort Belvoir,
VA 22060-6218

SUBJECT: Request Change in Distribution Statement

1. The U.S. Army Medical Research and Materiel Command has reexamined the need for the limitation assigned to technical reports. Request the limited distribution statement for reports on the enclosed list be changed to "Approved for public release; distribution unlimited." These reports should be released to the National Technical Information Service.

2. Point of contact for this request is Ms. Judy Pawlus at DSN 343-7322 or by e-mail at judy.pawlus@det.amedd.army.mil.

FOR THE COMMANDER:

Encl

PHYLLIS M. RINEHART
Deputy Chief of Staff for
Information Management

DAMD17-94-J-4413	ADB261602
DAMD17-96-1-6112	ADB233138
DAMD17-96-1-6112	ADB241664
DAMD17-96-1-6112	ADB259038
DAMD17-97-1-7084	ADB238008
DAMD17-97-1-7084	ADB251635
DAMD17-97-1-7084	ADB258430
DAMD17-98-1-8069	ADB259879
DAMD17-98-1-8069	ADB259953
DAMD17-97-C-7066	ADB242427
DAMD17-97-C-7066	ADB260252
DAMD17-97-1-7165	ADB249668
DAMD17-97-1-7165	ADB258879
DAMD17-97-1-7153	ADB248345
✓ DAMD17-97-1-7153	ADB258834
DAMD17-96-1-6102	ADB240188
DAMD17-96-1-6102	ADB257406
DAMD17-97-1-7080	ADB240660
DAMD17-97-1-7080	ADB252910
DAMD17-96-1-6295	ADB249407
DAMD17-96-1-6295	ADB259330
DAMD17-96-1-6284	ADB240578
DAMD17-96-1-6284	ADB259036
DAMD17-97-1-7140	ADB251634
DAMD17-97-1-7140	ADB259959
DAMD17-96-1-6066	ADB235510
DAMD17-96-1-6029	ADB259877
DAMD17-96-1-6020	ADB244256
DAMD17-96-1-6023	ADB231769
DAMD17-94-J-4475	ADB258846
DAMD17-99-1-9048	ADB258562
DAMD17-99-1-9035	ADB261532
DAMD17-98-C-8029	ADB261408
DAMD17-97-1-7299	ADB258750
DAMD17-97-1-7060	ADB257715
DAMD17-97-1-7009	ADB252283
DAMD17-96-1-6152	ADB228766
✓ DAMD17-96-1-6146	ADB253635
DAMD17-96-1-6098	ADB239338
DAMD17-94-J-4370	ADB235501
DAMD17-94-J-4360	ADB220023
DAMD17-94-J-4317	ADB222726
DAMD17-94-J-4055	ADB220035
DAMD17-94-J-4112	ADB222127
DAMD17-94-J-4391	ADB219964
DAMD17-94-J-4391	ADB233754